

DISCLOSURE OF POSSIBLE SIDE-EFFECTS

Semaglutide/Tirzepatide may cause tumors in the thyroid, including thyroid cancer. Watch for possible symptoms, such as a lump or swelling in the neck, hoarseness, trouble swallowing, or shortness of breath. If you have any of these symptoms, tell your healthcare provider.

- Do not use Semaglutide/Tirzepatide if you or any of your family have ever had a type of thyroid cancer called medullary thyroid carcinoma (MTC).
- Do not use Semaglutide/Tirzepatide if you have Multiple Endocrine Neoplasia syndrome type 2 (MEN 2).
- Do not use Semaglutide/Tirzepatide if you are allergic to it or any of the ingredients in Semaglutide/Tirzepatide.

Semaglutide/Tirzepatide may cause serious side effects, including:

Inflammation of the pancreas (pancreatitis). Stop using Semaglutide/Tirzepatide and call your healthcare provider right away if you have severe pain in your stomach area (abdomen) that will not go away, with or without vomiting. You may feel the pain from your abdomen to your back.

Low blood sugar (hypoglycemia). Your risk for getting low blood sugar may be higher if you use Semaglutide/Tirzepatide with another medicine that can cause low blood sugar, such as a sulfonylurea or insulin.

Signs and symptoms of low blood sugar may include dizziness or light-headedness, sweating, confusion or drowsiness, headache, blurred vision, slurred speech, shakiness, fast heartbeat, anxiety, irritability, or mood changes, hunger, weakness and feeling jittery.

Serious allergic reactions. Stop using Semaglutide/Tirzepatide and get medical help right away if you have any symptoms of a serious allergic reaction, including swelling of your face, lips, tongue or throat, problems breathing or swallowing, severe rash or itching, fainting or feeling dizzy, and very rapid heartbeat.

Kidney problems (kidney failure). In people who have kidney problems, diarrhea, nausea, and vomiting may cause a loss of fluids (dehydration), which may cause kidney problems to get worse. It is important for you to drink fluids to help reduce your chance of dehydration.

Severe stomach problems. Stomach problems, sometimes severe, have been reported in people who use Semaglutide/Tirzepatide. Tell your healthcare provider if you have stomach problems that are severe or will not go away.

Changes in vision. Tell your healthcare provider if you have changes in vision during treatment with Semaglutide/Tirzepatide.

Gallbladder problems. Gallbladder problems have happened in some people who use Semaglutide/Tirzepatide. Tell your healthcare provider right away if you get symptoms of gallbladder problems, which may include pain in your upper stomach (abdomen), fever, yellowing of skin or eyes (jaundice), and clay-colored stools.

Common side effects:

The most common side effects of Semaglutide/Tirzepatide include nausea, vomiting, decreased appetite, diarrhea, constipation, indigestion, and stomach (abdominal) pain.